

## On current risk factors associated with stroke types across gender and age in Southern Nigeria

A. Iduseri<sup>1, \*</sup> and P. Opoggen<sup>2</sup>

<sup>1,2</sup>*Department of Statistics, Faculty of Physical Sciences, University of Benin, Benin City, Nigeria*

*(Received: 04 April 2025; Accepted: 31 July 2025)*

**Abstract.** Stroke is the second leading cause of death worldwide, and remains one of the main causes of death in Nigeria, with its economic impact on the rise. Over the years, hospital-based studies in Nigeria often only report the traditional risk factors, with high blood pressure as the leading risk factor for cardiovascular disease, particularly stroke, in the country, without reporting any infectious risk factors. In this study, we will statistically assess all probable traditional and infectious risk factors associated with stroke. We aim to identify the current significant risk factors related to the prevalent type of stroke in Nigeria. This was a retrospective study conducted at a tertiary referral hospital, Federal Medical Centre (FMC), in Asaba, Delta State, Nigeria. We conducted a retrospective audit of medical records from 2019 to 2022 for adult stroke patients admitted to the medical ward. Data extracted included demographic characteristics, clinical variables, and relevant data. Identified risk factors associated with stroke types across gender and age were investigated using multinomial logistic regression. From the result, out of the recorded 907 stroke patients, 457 (50.39%) were female, while 450 (49.61%) were males. The age group (61-80) had more recorded cases totaling 426(47%). Hypertension, diabetes mellitus, heart disease, malaria, and pneumonia were the significant risk factors associated with the prevalent stroke type. Therefore, there is a need to step up measures aimed at increasing public awareness of emerging infectious risk factors associated with stroke and effective preventive measures. Prioritizing interventions directed towards cost-effective treatment of hospitalized stroke patients is imperative as this will hopefully improve the overall outcomes in resource-constrained settings such as Nigeria.

**Keywords:** Stroke types; traditional risk factors; infectious risk factors; multinomial logistic regression.

**Published by:** Department of Statistics, University of Benin, Nigeria

---

\*Corresponding author. Email:augustine.iduseri@uniben.edu

## 1. Introduction

A stroke is a sudden condition that leads to several related physical and cognitive impacts such as challenges with mobility, difficulty in communication, memory loss, depression, and sometimes being bedridden. It is an acute focal brain dysfunction originating from vessels and lasting 24 hours or longer. (Saengsuwan and Suangpho, 2019). According to the classification of the American Heart Association (AHA) report, the two major categories of strokes are ischemic and hemorrhagic stroke. Ischemic stroke, the most prevalent type, occurs from decreased or obstructed blood supply to the brain tissues, while hemorrhagic stroke occurs due to the burst of weakened blood vessels within the brain (Rink and Khanna, 2011). In Africa, the yearly incidence rate of stroke is about 316 cases per 100,000 person-years, while the prevalence rate is about 1,460 cases per 100,000 person-years (Akinyemi et al., 2021). Also, Africa has a 3-year fatality rate greater than 80%, with stroke-related deaths being about 5.5% to 11% of mortality (Akinyemi et al., 2021). Current evidence indicates that Africa could have up to 2-3-fold greater rates of stroke incidence and higher stroke prevalence than Western Europe and the USA (Tamer et al., 2020). The increase in stroke burden in Africa is being driven mostly by growing trends in the prevalence of modifiable risk factors (which include smoking, hypertension, dyslipidemia, hyperlipidemia, diabetes, poor diet, abdominal obesity, physical inactivity, atrial fibrillation, high salt intake), and also regulated by non-modifiable risk factors (such as race, sex, age, geographic locations, and heredity) (Onwuakagba et al., 2023). Nigeria, similar to many other low-income and middle-income countries in Sub-Saharan Africa, has been experiencing a significant increase in stroke burden with an estimated yearly incidence rate of 1.14 per 1000 persons/year (Moghtaderi and Alavi-Naini, 2012). The increasing number of cardiovascular diseases (CVD), especially stroke, is partly attributed to the growing presence of both modifiable and non-modifiable risk factors. Additionally, urbanization and the adoption of Western lifestyles, characterized by sedentary lifestyles, harmful habits like tobacco and alcohol consumption, and high fat/cholesterol diets, also contribute to this rise. (Omigbile et al., 2023; Ike and Onyema, 2020). Stroke remains a leading cause of physical or mental impairment, and the second most common cause of death globally and is associated with substantial economic costs for the sufferers, and families (Onwuakagba et al., 2023; Teh et al., 2018). Surprisingly, the consequences on patients and their families are often unexplainable and can even be referred to as tragic because of the enormous psychological, social, and financial burden associated with stroke.

Several hospital-based studies in Nigeria have documented cardiovascular disease, particularly stroke, as the most common cause of neurological admissions and medical coma (Talabi, 2003; Ojini and Danesi, 2003; Ogun et al., 2005; Ekenze et al., 2010; Eze and Kalu, 2014; Kayode-Iyasere et al., 2019; Arabambi et al., 2021; Osuegbu et al., 2021; Nwazor et al., 2024). The common risk factors identified by these hospital-based studies in Nigeria are high blood pressure, diabetes, consuming tobacco products, physical inactivity, unhealthy diet, and aging. While traditional cardiovascular risk factors had been reported as common causes for most strokes in Nigeria, Sub-Saharan Africa and other parts of the globe, infectious pathogens have also been reported to increase the risk, and in

some cases, have a direct causal role (Moghtaderi and Alavi-Naini, 2012; Jennifer and Fugate, 2020; Hameed et al., 2024). Important examples of infectious pathogens or infectious diseases of stroke (i.e. parasitic, bacterial, viral, or fungal infections) that can result in stroke include cerebral malaria (Carod-Artal, 2007; Leopoldino et al., 1999; Periyasamy et al., 2023), tuberculous meningitis (Wasay et al., 2018), pneumonia (Ingeman et al., 2011; Kim et al., 2017; Grossmann et al., 2021; Watila et al., 2019; Olajide et al., 2021), HIV/AIDS (Grau et al., 2010; Thakur et al., 2016), cysticercosis (Alarcón et al., 1992a; Alarcón et al., 1992b), syphilis (Jennifer and Fugate, 2020), chronic hepatitis infection (Cojocarú et al., 2005; Sung et al., 2007), Chagas disease (Lage et al., 2022), and most recently, coronavirus disease 2019 (COVID-19) (Khan et al., 2023). Notably, infectious risk factors or tropical diseases such as malaria, pneumonia, and HIV/AIDS that are not frequently among the primary risk factors in developed countries, are endemic in Nigeria. In 2021, Nigeria recorded an estimated 68 million cases and 194,000 deaths from malaria (WHO, 2022). Some case presentations on stroke have indicated a link with the most severe complication of malaria caused by *Plasmodium falciparum*, also known as cerebral malaria (CM), which could be accountable for up to 10% of strokes in endemic regions like Nigeria (Carod-Artal, 2007; Leopoldino et al., 1999; Periyasamy et al., 2023). Similarly, Warren-Gash et al. (2018) in their study found that individuals with pneumonia could have an increased risk of heart attack for up to one week after infection, and stroke for up to 28 days. In Nigeria, community-acquired pneumonia (CAP) accounts for 2.5% to 5.7% of medical admissions and between 15.3% and 24.9% of respiratory admissions in tertiary health facilities. Previous surveys have shown that mortality rates among admitted patients with CAP range from 7.4% to 26% in the country. Besides the infectious risk factors, environmental risk factors are also emerging as non-traditional risk factors associated with stroke (Hameed et al., 2024). Important examples include air pollution (Wellenius et al., 2012; Feigin et al., 2016), climate change (Chu et al., 2018; Atwoli et al., 2021), and high altitude (Jaillard et al., 1995; Lu et al., 2020; Liu et al., 2021). Air pollution, which is more noticeable in developing countries, is a recognized risk factor for cardiovascular disease and reportedly accounts for 14% of all stroke-associated mortality (Verhoeven et al., 2021). Currently, Nigeria like many other Sub-Saharan African (SSA) countries is generally going through health transition. This transition is characterized by increasing urbanization and lifestyle changes, resulting in a more prevalence of cardiovascular diseases, as well as infectious and environmental risk factors associated with stroke.

Many hospital-based studies have confirmed cardiovascular disease, particularly stroke, as the most common cause of neurological admissions and medical coma in Nigeria. However, these studies often only report the traditional risk factors, with high blood pressure as the primary risk factor for stroke in Nigeria, without reporting any infectious risk factors. Against this background, this study employed multinomial logistic regression to ascertain if there are significant relationships between any infectious risk factors and stroke types in Southern Nigeria, relying on the evidence from retrospective hospital data. We aim to identify the current significant risk factors associated with the prevalent type of stroke in Nigeria. The specific objectives of this study are: (1) To evaluate the

association between different age groups and various types of strokes. (2) To evaluate the association of sex and various types of strokes. (3) To statistically ascertain the current traditional and infectious risk factors associated with the types of strokes in Nigeria. To achieve these objectives, the remainder of the paper is structured as follows: Section 2 describes the research materials and methods employed in the study. Section 3 discusses the results of our inquiry, while Section 4 presents our concluding remarks.

## **2. Materials and Method**

### **2.1 Study Design and Data Sources**

This hospital-based retrospective study was conducted at the Federal Medical Centre (FMC) in Asaba. Asaba town is the capital of Delta State in Southern Nigeria located at the western bank of the Niger River. Asaba, with a population of 149,603 as of the 2006 census, is close to Onitsha, Owerri, and Enugu, which are known major cities in Southeastern Nigeria. This makes FMC, Asaba a referral center for major cities in both the South-South and South-East regions of Nigeria. The target population comprises patients having cerebrovascular accident (CVA) or presenting with acute stroke at FMC, Asaba, between August 2019 and September 2022. Medical records of 907 stroke patients were retrieved, and relevant information was extracted using a structured questionnaire. The questionnaire contained relevant information such as a unique case identifier, age, sex, duration to presentation, duration to outcome, length of hospitalization, clinical indications (stroke type), risk factors (both vascular and infectious risk factors), and access to neuroimaging. Eight categories of stroke type were identified as reported in their admission records. These include Transient Ischemic Stroke (TIS), Ischemic Stroke (IS), Acute Stroke (AS), Hemorrhagic Stroke (HS), Left Hemispheric Stroke (LHS), Left Hemispheric Ischemic Stroke (LHIS), Right Hemispheric Stroke (RHS), and Right Hemispheric Ischemic Stroke (RHIS).

Ethical clearance for this study was obtained from the Research and Ethical Committee, Federal Medical Centre, Asaba. The ethical reference number was FMC/ASB/A81 VOL XII/268. The permission to collect data from the patient's medical records was also obtained from the Department of Medical Records. During the research, we ensured anonymity and confidentiality were upheld to the highest standard.

### **2.2 Statistical Analysis**

The obtained data was inputted into IBM Statistical Package for the Social Sciences (SPSS) Statistics Version 26 Data Editor for the analysis. Before starting the analysis, we checked for missing values and used the SPSS multiple imputation technique to handle them. SPSS descriptive statistics; frequencies (percentages) for categorical variables were used to describe the data distributions of some sociodemographic characteristics and other variables of the sample data for presentation. We then applied SPSS multinomial logistic regression to assess the association between different age groups and sex with the types of strokes,

and to identify the current significant risk factors associated with the types of strokes. Because we are analyzing a set of categorical variables, and one is a “response” while the others are predictors, we adopted the multinomial logistic regression to achieve our objectives. For a multinomial logistic regression, the predicted probability of having success in the  $k^{th}$  category and a given input  $x$  is given in Equation 2.1a

$$P(Y_j = k | X_{1,i}, \dots, X_{p,i}) = \frac{\exp(\beta_k x_i)}{1 + \sum_{j=0}^{K-1} \exp(\beta_j x_i)} \quad (2.1a)$$

To ensure identifiability (a unique solution), one class is chosen as the reference class, and its coefficients are set to zero. Hence, the predicted probability becomes

$$P(Y_j = k | X_{1,i}, \dots, X_{p,i}) = \frac{\exp(\beta_k x_i)}{1 + \sum_{j=0}^{K-1} \exp(\beta_j x_i)} \text{ for } k = 1, \dots, K - 1 \quad (2.1b)$$

The  $k^{th}$  logistic regression model for the variables applicable in this study is given by

$$\ln \left[ \frac{P(Y_j = k | X_{1,i}, \dots, X_{8,i})}{P(Y_j = 0 | X_{1,i}, \dots, X_{8,i})} \right] = \beta_{0,k} + \beta_{1,k} x_1 + \beta_{2,k} x_2 + \dots + \beta_{8,k} x_8$$

$$\text{for } k = 1, \dots, 5 \quad (2.2)$$

where  $Y_j$  is the value of the multinomial response variable for the  $i^{th}$  unit,  $X_{1,i}, \dots, X_{8,i}$  are the eight categories of stroke type or predictor variables that were identified as reported in their admission records,  $\beta_0, \beta_1, \dots, \beta_8$  are the coefficients of the eight categories of stroke type, and  $K^{th}$  category (which is the sixth category of stroke type) is the reference category. Each coefficient  $\beta_i$  in the  $k^{th}$  category of stroke type represents the log-odds change of being in category  $k$  relative to the reference class, for a one-unit increase in the predictor  $X_i$ . In the analysis of multinomial logistic regression outputs, the probabilities obtained from Equation 2.1b will enable us to determine classification, while the coefficients that will be obtained from Equation 2.2 will guide us in understanding the influence or assessing the association of risk factors with the prevalent type of stroke, which is the primary focus of this study.

### 3. Results and Discussion

#### 3.1 Differences in stroke type prevalence across patients' sex

The results in Table 1 showed that there are 457(50.39%) female patients with stroke, and 450 (49.61%) male patients with stroke. Our observed trend of higher stroke prevalence among women aligns with a previous study conducted

at FMC, Asaba, Delta State, Nigeria (Ezunu et al., 2023). Upon reviewing Table 1, it is evident that ischemic stroke was more common in females (46.6%), followed by hemorrhagic stroke (11.6%), and right hemispheric stroke (11.2%). Ischemic stroke (55.1%) was also the most common for males, followed by right hemispheric stroke (12.2%). In the present study, it was found that 461 (50.83%) patients had ischemic stroke, the highest incidence among the patients. This finding is consistent with several other studies conducted in different African countries. The similarity in the genetic makeup of different race/ethnicity may account for this observation. However, only 39 (4.30%) of the patients had acute stroke, which is the lowest type of stroke recorded in this study.

Table 1: Stroke types across sex and their proportions

| Stroke Types                           | No. of Patients | Female      | Males       |
|----------------------------------------|-----------------|-------------|-------------|
| Transient ischemic stroke (TIS)        | 46(5.07%)       | 30(6.6%)    | 16(3.6%)    |
| Ischemic stroke (IS)                   | 461(50.83%)     | 213(46.6%)  | 248(55.1%)  |
| Acute stroke (AS)                      | 39(4.30%)       | 11(2.4%)    | 28(6.2%)    |
| Hemorrhagic stroke (HS)                | 84(9.26%)       | 53(11.6%)   | 31(6.9%)    |
| Left hemispheric stroke (LHS)          | 47(5.18%)       | 28(6.1%)    | 19(4.2%)    |
| Left hemispheric ischemic stroke(LHIS) | 48(5.29%)       | 34(7.4%)    | 14(3.1%)    |
| Right hemispheric stroke (RHS)         | 106(11.69%)     | 51(11.2%)   | 55(12.2%)   |
| Left hemispheric ischemic stroke(LHIS) | 76(8.38%)       | 37(8.1%)    | 39(8.7%)    |
|                                        |                 | 457(50.39%) | 450(49.61%) |

This shows that ischemic stroke was more probable to cause death in the studied population. This finding is consistent with previous conducted studies (Jowi and Mativo, 2008; Zhang et al., 2017; Mulugeta et al., 2020; Arabambi et al., 2021). On the other hand, in other studies in Nigeria and the rest of the world, hemorrhagic stroke was the most common subtype (Gedefa et al., 2017; Zewdie et al., 2018; Maskey et al., 2011; Osuegbu et al., 2021). This difference may be attributed to underlying differences in risk factors (Zhang et al., 2017).

### 3.2 Differences in stroke type prevalence across patients' different age groups

Table 2 shows the distribution of stroke patients by age group. There are 19 patients with stroke in the age group of 20 or younger, 45 patients in the age group 21-40, 286 patients in the age group 41-60, 426 patients in the age group 61-80, and 131 in the age group 81-100. This data indicates that older age groups have a higher prevalence of stroke. The research conducted by Teh et al. in 2018 and Osuegbu et al. in 2021 indicates that as individuals get older, they are more likely to experience a stroke. This could be attributed to the physiological changes that occur in the elderly, as well as the greater complexity and severity of the condition, which tend to increase with age.

Table 2: Stroke types across different age group and their proportions

| Stroke Types | 20        | 21-40     | 41-60       | 61-80       | 81-100      |
|--------------|-----------|-----------|-------------|-------------|-------------|
| TIS          | 0(0.0%)   | 2(4.4%)   | 12(4.2%)    | 25(5.9%)    | 7(5.3%)     |
| IS           | 7(36.8%)  | 19(42.2%) | 143(50.0%)  | 241(56.6%)  | 51(38.9%)   |
| AS           | 2(10.5%)  | 0(0.0%)   | 1(0.3%)     | 20(4.7%)    | 16(12.2%)   |
| HS           | 6(31.6%)  | 6(13.3%)  | 40(14.0%)   | 23(5.4%)    | 9(6.9%)     |
| LHS          | 0(0.0%)   | 5(11.1%)  | 20(7.0%)    | 18(4.2%)    | 4(3.1%)     |
| LHIS         | 2(10.5%)  | 2(4.4%)   | 14(4.9%)    | 20(4.7%)    | 10(7.6%)    |
| RHS          | 0(0.0%)   | 4(8.9%)   | 34(11.9%)   | 40(9.4%)    | 28(21.4%)   |
| RHIS         | 2(10.5%)  | 7(15.6%)  | 22(7.7%)    | 39(9.2%)    | 6(4.6%)     |
|              | 19(2.09%) | 45(4.96%) | 286(31.54%) | 426(46.97%) | 131(14.44%) |

Consequently, older individuals may require more inpatient health services due to these factors. A cursory look at Table 2 shows that the distribution of stroke types in the age group 61-80 is relatively higher in terms of incidence than in any other age group. This means that older adults dominated the stroke incidence and were more likely to die from a stroke, irrespective of the type of stroke. However, the prevalence of ischemic stroke was higher across the age groups. This again shows that ischemic stroke was more likely to cause death in the studied population.

### 3.3 Predominant risk factors across patients' sex

Table 3 presents the distribution of the predominant risk factors among the studied population. Out of all the potential risk factors listed in the structured questionnaire used for data collection, including cardiovascular diseases and infectious risk factors, only the six risk factors shown in Table 3 were predominantly documented as clinical diagnoses and/or laboratory diagnoses in the medical records of stroke patients.

Table 3: Predominant risk factors across sex and their proportions

| Risk Factors      | No. of Patients | Male       | Females     |
|-------------------|-----------------|------------|-------------|
| Hypertension      | 81              | 34(41.98%) | 47(58.02%)  |
| Diabetes Mellitus | 56              | 16(28.75%) | 40(71.43%)  |
| Malaria           | 22              | 14(63.64%) | 8 (36.6%)   |
| Pneumonia         | 20              | 9 (45.00%) | 11 (55.00%) |
| Heart Disease     | 24              | 5 (20.83%) | 19 (79.17%) |
| Chronic Anemia    | 5               | 5(100.00%) | 0 (0.00%)   |

While other risk factors listed in the structured questionnaire had only one or two recorded cases in the medical records of the studied population.

In Table 3, hypertension and diabetes mellitus, were found to have the highest recorded risk factors for stroke patients in the studied population, which is consistent with previous studies (Mulugeta et al., 2020; Ojo et al., 2024). The results in Table 3 also indicate that heart disease, malaria, and pneumonia were the second-highest recorded infectious risk factors. Malaria was observed more frequently in males (63.64%) while pneumonia was more commonly observed in females (55.00%). Previous studies have linked severe complications of malaria caused by *Plasmodium falciparum* and pneumonia as possible infec-

tious risk factors associated with stroke (Carod-Artal, 2007; Leopoldino et al., 1999; Warren-Gash et al., 2018; Periyasamy et al., 2023).

To determine if the predominant risk factors listed in Table 3 are significantly associated with different types of strokes, we used multinomial logistic regression. In this analysis, the risk factors were used as the multinomial response variable, while the stroke types were used as the predictor variables.

Table 4: Multinomial logistic regression model for stroke types and risk factors

| Risk factor              | $\beta$        | S.E       | Wald     | df | Sig   | Exp( $\beta$ ) | [95% C.I]                |
|--------------------------|----------------|-----------|----------|----|-------|----------------|--------------------------|
| <b>Hypertension</b>      |                |           |          |    |       |                |                          |
| Intercept                | 19.735         | 0.880     | 503.013  | 1  | 0.000 |                |                          |
| TIS                      | -18.348        | 1.423     | 166.314  | 1  | 0.000 | 1.075E-8       | [6.613E-10; 1.748E-7]    |
| IS                       | -17.687        | 0.746     | 562.131  | 1  | 0.000 | 2.083E-8       | [4.827E-9 ; 8.988E-8]    |
| AS                       | 0.159          | 0.867     | 0.033    | 1  | 0.856 | 1.173          | [0.211; 6.523]           |
| HS                       | 0.957          | 11763.256 | 0.000    | 1  | 1.000 | 2.603          | [0.000; . <sup>b</sup> ] |
| LHS                      | 0.598          | 13002.555 | 0.000    | 1  | 1.000 | 1.818          | [0.000; . <sup>b</sup> ] |
| LHIS                     | 0.038          | 7431.315  | 0.000    | 1  | 1.000 | 1.039          | [0.000; . <sup>b</sup> ] |
| RHS                      | 0.673          | 12079.281 | 0.000    | 1  | 1.000 | 1.961          | [0.000; . <sup>b</sup> ] |
| RHIS                     | 0 <sup>c</sup> | .         | .        | 0  | .     | .              | .                        |
| <b>Diabetes mellitus</b> |                |           |          |    |       |                |                          |
| Intercept                | 19.378         | 0.904     | 459.557  | 1  | 0.000 |                |                          |
| TIS                      | -18.279        | 1.466     | 155.379  | 1  | 0.000 | 1.152E-8       | [6.504E-10; 2.040E-7]    |
| IS                       | -17.720        | 0.784     | 510.782  | 1  | 0.000 | 2.016E-8       | [4.336E-9; 9.372E-8]     |
| AS                       | 0.110          | 0.923     | 0.014    | 1  | 0.905 | 1.117          | [0.183; 6.812]           |
| HS                       | 0.466          | 11763.56  | 0.000    | 1  | 1.000 | 1.594          | [0.000; . <sup>b</sup> ] |
| LHS                      | 0.954          | 13002.55  | 0.000    | 1  | 1.000 | 2.597          | [0.000; . <sup>b</sup> ] |
| LHIS                     | 0.395          | 7431.315  | 0.000    | 1  | 1.000 | 1.484          | [0.000; . <sup>b</sup> ] |
| RHS                      | 0.807          | 12079.218 | 0.000    | 1  | 1.000 | 2.241          | [0.000; . <sup>b</sup> ] |
| RHIS                     | 0 <sup>c</sup> | .         | .        | 0  | .     | .              | .                        |
| <b>Malaria</b>           |                |           |          |    |       |                |                          |
| Intercept                | 18.818         | 0.961     | 383.155  | 1  | 0.000 |                |                          |
| TIS                      | -18.818        | 1.710     | 121.101  | 1  | 0.000 | 6.719E-9       | [2.354E-10; 1.918E-7]    |
| IS                       | -17.566        | 0.864     | 413.786  | 1  | 0.000 | 2.352E-8       | [4.329E-9; 1.278E-7]     |
| AS                       | -1.409         | 1.354     | 1.083    | 1  | 0.298 | 0.244          | [0.017; 3.471]           |
| HS                       | -17.337        | 13031.610 | 0.000    | 1  | 0.999 | 2.956E-8       | [0.000; . <sup>b</sup> ] |
| LHS                      | -17.337        | 14404.533 | 0.000    | 1  | 0.999 | 2.956E-8       | [0.000; . <sup>b</sup> ] |
| LHIS                     | -0.298         | 7431.315  | 0.000    | 1  | 1.000 | 0.742          | [0.000; . <sup>b</sup> ] |
| RHS                      | -17.337        | 13381.713 | 0.000    | 1  | 0.999 | 2.956E-8       | [0.000; . <sup>b</sup> ] |
| RHIS                     | 0 <sup>c</sup> | .         | .        | 0  | .     | .              | .                        |
| <b>Pneumonia</b>         |                |           |          |    |       |                |                          |
| Intercept                | 18.818         | 0.961     | 383.155  | 1  | 0.000 |                |                          |
| TIS                      | -36.244        | 6079.164  | 0.000    | 1  | 0.995 | 1.818E-16      | [0.000; . <sup>b</sup> ] |
| IS                       | -18.007        | 0.886     | 412.891  | 1  | 0.000 | 1.512E-8       | [2.662E-9; 8.587E-8]     |
| AS                       | -0.023         | 1.041     | 0.000    | 1  | 0.982 | 0.977          | [0.127; 7.515]           |
| HS                       | -17.483        | 13220.58  | 0.000    | 1  | 0.999 | 2.553E-8       | [0.000; . <sup>b</sup> ] |
| LHS                      | -17.483        | 14613.08  | 0.000    | 1  | 0.999 | 2.553E-8       | [0.000; . <sup>b</sup> ] |
| LHIS                     | -0.298         | 7431.315  | 0.000    | 1  | 1.000 | 0.742          | [0.000; . <sup>b</sup> ] |
| RHS                      | -17.483        | 13575.942 | 0.000    | 1  | 0.999 | 2.553E-8       | [0.000; . <sup>b</sup> ] |
| RHIS                     | 0 <sup>c</sup> | .         | .        | 0  | .     | .              | .                        |
| <b>Heart disease</b>     |                |           |          |    |       |                |                          |
| Intercept                | 18.531         | 0.584     | 1007.259 | 1  | 0.000 |                |                          |
| TIS                      | -16.921        | 1.241     | 185.817  | 1  | 0.000 | 4.479E-8       | [3.932E-9; 5.103E-7]     |
| IS                       | -17.519        | 0.000     | .        | 1  | .     | 2.464E-8       | [2.464E-8; 2.464E-8]     |
| AS                       | -0.265         | 0.000     | .        | 1  | .     | 1.303E-8       | [1.303; 1.303]           |
| HS                       | -16.962        | 12930.666 | 0.000    | 1  | 0.999 | 4.300E-8       | [0.000; . <sup>b</sup> ] |
| LHS                      | -16.962        | 14292.954 | 0.000    | 1  | 0.999 | 4.300E-8       | [0.000; . <sup>b</sup> ] |
| LHIS                     | -0.704         | 7431.315  | 0.000    | 1  | 1.000 | 0.495          | [0.000; . <sup>b</sup> ] |
| RHS                      | -16.962        | 13278.057 | 0.000    | 1  | 0.999 | 4.300E-8       | [0.000; . <sup>b</sup> ] |
| RHIS                     | 0 <sup>c</sup> | .         | .        | 0  | .     | .              | .                        |

a. The reference category is: chronic anemia b. Floating point overflow occurred while computing this statistic. Its value is therefore set to system missing. c. This parameter is set to zero because it is redundant.

Table 4 displays the SPSS output of the multinomial logistic regression analysis

for the stroke types and risk factors as presented in Equation (2.2). A cursory look at Table 4 reveals that transient ischemic stroke (TIS) and ischemic stroke (IS) have a significant relationship with hypertension ( $p=0.000$ ), diabetes mellitus ( $p=0.000$ ), and malaria ( $p=0.000$ ), respectively. Table 4 also reveals that IS has a significant relationship with pneumonia ( $p=0.000$ ), and TIS has a significant relationship with heart disease ( $p=0.000$ ). Column 2 of Table 4 presents the Wald statistic for TIS and IS, corresponding to their coefficient values for the five predominant risk factors. The observed substantially large Wald statistic presents strong evidence that TIS and IS coefficient values are significantly different from zero. Additionally, in columns 2 and 3 of Table 4, we also observed that the standard errors (SE) are much smaller than the absolute values of the coefficients for TIS and IS for the five predominant risk factors, which is typically good, and a confirmation that TIS and IS have a very significant relationship with the predominant risk factors.

This result not only confirms our earlier assertion on Table 3 results, but further reveals that all the identified predominant risk factors have a significant relationship with ischemic stroke. This suggests that among all stroke types, ischemic stroke (whether TIS or IS) has the highest prevalence in this study, which is consistent with previous studies (Mulugeta et al., 2020; Osuegbu et al., 2021). In addition to the commonest risk factors such as hypertension, diabetes mellitus, and heart disease identified in this study, the significant relationship of malaria and pneumonia with ischemic stroke in Table 4 suggests the two infectious risk factors as significant emerging risk factors for stroke in Nigeria. This finding is corroborated by previous studies and presentations outside Nigeria that have linked severe complications of malaria caused by *plasmodium falciparum* and pneumonia as possible infectious risk factors associated with stroke, and others that even suggested they could have a direct causal role (Carod-Artal, 2007; Leopoldino et al., 1999; Warren-Gash et al., 2018; Periyasamy et al., 2023).

#### **4. Conclusion**

This study found that ischemic stroke was a common indication for medical admissions, in particular for the age group 61-80, with hypertension and diabetes mellitus as the leading risk factors, followed by heart disease, malaria, and pneumonia. Therefore, there is a need to step up measures aimed at increasing public awareness of emerging infectious risk factors (such as malaria and pneumonia) associated with stroke and providing effective preventive measures. Prioritizing interventions directed towards cost-effective treatment of hospitalized stroke patients is imperative, as this will hopefully improve the overall outcomes in resource-constrained settings such as Nigeria.

This study, being a retrospective review of records, had some limitations. Many of the case files had substantial missing data and could not be included in the data analysis. This low yield of data may distort the accurate picture of stroke presentations at the hospital. Also, the measured risk factors were likely under-reported as they were absent in some of the case files reviewed. A multi-centered prospective and retrospective, registry-based study that will provide more representative and reliable data regarding stroke types with current associated risk factors in Nigeria will, without doubt, help establish this study result.

## References

- Akinyemi, R.O., Ovbiagele, B., Adeniji, O.A. et al. (2021). Stroke in Africa: profile, progress, prospects and priorities. *Nat Rev Neurol*, 17, 634–656. <https://doi.org/10.1038/s41582-021-00542-4>
- Alarcón, F., Hidalgo, F., Moncayo, J., Viñán, I., Dueñas, G. (1992a). Cerebral cysticercosis and stroke. *Stroke*, 23: 224–8. doi: 10.1161/01.STR.23.2.224. PubMed PMID: 1561652.
- Alarcón, F., Vanormelingen, K., Moncayo, J., Viñán, I. (1992b). Cerebral cysticercosis as a risk factor for stroke in young and middle-aged people. *Stroke*, 23, 1563–5. doi: 10.1161/01.STR.23.11.1563. PubMed PMID: 1440703.
- Arabambi, B., Oshinaike, O., Akilo, O.O., Yusuf, Y., Ogun, S.A. (2021). Pattern, risk factors, and outcome of acute stroke in a Nigerian university teaching hospital: a 1-year review. *Nigerian Journal of Medicine*, 30(3), 252–258. DOI: 10.4103/NJM.NJM-8-21
- Atwoli, L., Baqui, A.H., Benfield, T., Bosurgi, R., Godlee, F., Hancocks, S., Horton, R.C., Laybourn-Langton, L., Monteiro, C.A., Normanet, I., et al. (2021). Call for emergency action to limit global temperature increases, restore biodiversity, and protect health. *BMJ Nutr. Prev. Health*, 4, 362–364.
- Carod-Artal, F.J. (2007). Strokes caused by infection in the tropics. *Rev Neurol*, 44:755–63. PubMed PMID: 17583870. [PubMed]
- Chu, S.Y.; Cox, M., Fonarow, G.C., Smith, E.E., Schwamm, L., Bhatt, D.L., Matsouaka, R.A., Xian, Y., Sheth, K.N. (2018). Temperature and precipitation associate with ischemic stroke outcomes in the United States. *J. Am. Heart Assoc.*, 7, e010020.
- Cojocar, I.M., Cojocar, M., Iacob, S.A., Burcin, C. (2005). Cerebral ischemic attack secondary to hepatitis C virus infection. *Rom. J. Intern. Med*, 43, 255–260.
- Ekenze, O., Onwuekwe, I. and Ezeala, B. (2010). Profile of neurological admissions at the university of Nigeria teaching hospital Enugu. *Niger. J. Med*, 19, 419–422.
- Eze, C. O. and Kalu, U. A. (2014). Pattern of neurological admissions in the topics: experience at Abakaliki South-Eastern Nigeria. *Niger. J. Med*, 23, 302–305.
- Ezunu, O.E., Akpekpe, J., Yusuf, Y., Ogunniyi, A., Ige, F., Ezunu, N.E., Ogbutor, U. G., Guobadia, E. K. (2023). Pattern and distribution of neurological disorders among patients that attended Federal Medical Centre, Asaba, Nigeria between 2015-2019. *International Journal of Research and Innovation in Social Science (IJRISS)*, ISSN No. 2454-6186
- Feigin, V.L., Roth, G.A., Naghavi, M., Parmar, P., Krishnamurthi, R., Chugh, S., Mensah, G.A., Norrving, B., Shiue, I., Ng, M., et al. (2016). Global burden of stroke and risk factors in 188 countries, during 1990–2013: a systematic analysis for the global burden of disease study 2013. *Lancet Neurol.*, 15, 913–924.
- Gedefa, B., Menna, T., Berhe, T., Abera, H. (2017). Assessment of risk factors and treatment outcome of stroke admissions at St. Paul's teaching hospital, Addis Ababa, Ethiopia. *J Neurol and Neurophysiol*, 8(3), 1–6.
- Grau, A.J., Urbanek, C., Palm, F. (2010). Common Infections, and the Risk of Stroke. *Nat. Rev. Neurol*, 6, 681–694.
- Grossmann, I., Rodriguez, K., Soni, M., Joshi, P.K., Patel, S.C., Shreya, D., Zamora, D.I., Patel, G.S., Sange, I. (2021) Stroke and pneumonia: mechanisms, risk factors, management, and prevention. *Cureus*, 13(11), e19912. doi:10.7759/cureus.19912
- Hameed, S., Karim, N., Wasay, M., and Venketasubramania, N. (2024). Emerging stroke risk factors: a focus on infectious and environmental determinants. *J. Cardiovasc. Dev. Dis.*, 11(1): 19; <https://doi.org/10.3390/jcdd11010019>
- Ike, S. and Onyema, C. (2020). Cardiovascular diseases in Nigeria: what has happened in the past 20 years? *Nig J Cardiol*, (17), 21–26.
- Ingeman, A., Andersen, G., Hundborg, H.H., Svendsen, M.L., Johnsen, S.P. (2011). In-hospital medical complications, length of stay, and mortality among stroke unit patients. *Stroke*, 42,3214–3218. [PubMed] [Google Scholar]
- Jaillard, A.S., Hommel, M., Mazetti, P. (1995). Prevalence of stroke at high altitude (3380 m) in Cuzco, a town of Peru. a population-based study. *Stroke*, 26, 562–568.
- Jennifer E. and Fugate, D.O. (2020). Infectious causes of stroke. *Practical Neurology*, <https://practicalneurology.com/articles/2020-jan/infectious-causes-of-stroke>

- Jowi, J., and Mativo, P. (2008). Pathological sub-types, risk factors and outcome of stroke at the Nairobi Hospital, Kenya. *East Afr Med J*, 85(12), 572–81.
- Kayode-Iyasere, E. O., Obasohan, A. O. and Odiase, F. E. (2019). Medical coma in a secondary health centre in Benin City, Nigeria: a 3-year review. *Port Harcourt Med. J*, 13, 58–62.
- Khan, M., Hameed, S., Soomro, B.A., Mairaj, S., Malik, A., Farooq, S., Rukn, S.A., Wasay, M. (2023). COVID-19 independently predicts poor outcomes in acute ischemic stroke-insights from a multicenter study from Pakistan and United Arab Emirates. *J. Stroke Cerebrovasc. Dis*, 32, 106903.
- Kim, B.R., Lee, J., Sohn, M.K., Kim, D.Y., Lee, S.G., Shin, Y.I., Oh, G.J., Lee, Y.S., Joo, M.C., Han, E.Y., and Kim, Y.H., (2017). Risk factors and functional impact of medical complications in stroke. *Ann Rehabil Med*, 41,753–760. [PMC free article] [PubMed] [Google Scholar]
- Lage, T.A.R., Tupinambás, J.T., Pádua, L.B., Ferreira, M.O., Ferreira, A.C., Teixeira, A.L., Nunes, M.C.P. (2022). Stroke in chagas disease: from pathophysiology to clinical practice. *Rev. Soc. Bras. Med. Trop*, 55, e0575.
- Leopoldino, J. F. S., Fukujima, M. M., and Gabbai, A. A. (1999). Malaria and stroke: case report, *Arquivos De Neuro-psiquiatria*, 57(4), 1024–1026. <https://doi.org/10.1590/S0004-282X1999000600021>
- Liu, M., Yan, M., Guo, Y., Xie, Z., Li, R., Li, J., Ren, C., Ji, X., Guo, X. (2021). Acute ischemic stroke at high altitudes in China: early onset and severe manifestations. *Cells*, 10, 809.
- Lu, Y., Zhuoga, C., Jin, H., Zhu, F., Zhao, Y., Ding, Z., He, S., Du, A., Xu, J., Luo, J., et al. (2020). Characteristics of acute ischemic stroke in hospitalized patients in Tibet: a retrospective comparative study. *BMC Neurol*, 20, 380.
- Maskey, A., Parajuli, M., Kohli, S. C. (2011). A study of risk factors of stroke in patients admitted in Manipal Teaching Hospital, Pokhara. *Kathmandu Univ Med J (KUMJ)*, 9(36), 244–7.
- Moghtaderi, A. and Alavi-Naini, R. (2012). Infective causes of stroke in tropical regions. *Iran J Med Sci*, 37(3), 150–158. PMID: 23115446.
- Mulugeta, H., Yehuala, A., Haile, D., Mekonnen, N., Dessie, G., Kassa, G. M., Kassa, Z. S., and Habtewold, T. D. (2020). Magnitude, risk factors and outcomes of stroke at Debre Markos Referral Hospital, Northwest Ethiopia: a retrospective observational study. *Egypt J Neurol Psychiatry Neurosurg* 56, 41. <https://doi.org/10.1186/s41983-020-00173-4>
- Nwazor, E., Obi, P., Madueke, O., Ajuonuma, B., and Chukwuocha, I. (2024). A 2-year review of stroke admissions and short term out-come predictors in a teaching hospital, Southeast, Nigeria. *Nigerian Medical Journal*, 65(2), 185–194. <https://doi.org/10.60787/nmj-v65i2-442>
- Ogun, S., Ojini, F., Ogungbo, B., Kolapo, K. and Danesi, M. (2005). Stroke in South West Nigeria: a 10-year review. *Stroke*, 36, 1120–1122.
- Ojini, F. and Danesi, M. (2003). The Pattern of neurological admissions at the Lagos university teaching hospital. *Niger. J. Clin. Pract*, 6, 38–41.
- Ojo, A.E., Ojji, D.B., Grobbee, D.E., Huffman, M.D., Peters, S.A.E. (2024). The burden of cardiovascular disease attributable to hypertension in Nigeria: a modelling study using summary-level data. *Global Heart*, 19(1), 50. DOI: <https://doi.org/10.5334/gh.1332>
- Olajide, O.O., Adebayo, P.B., Taiwo, F.T., Owolabi, M.O., and Ogunniyi, A. (2021). The contribution of dysphagia to stroke morbidity and mortality among Nigerians, *Annals of Health Research*, 7(2), 118-131. doi:10.30442/ahr.0702-04-123.
- Omigbile, O., Oni, A., Abe, E., Lawal, E., Oyasope, B., and Mayaki, T. (2023). Prevalence and risk factors of cardiovascular diseases among the Nigerian population: a new trend among adolescents and youths. In book: novel pathogenesis and treatments for cardiovascular disease. Intech open, DOI: 10.5772/intechopen.108180
- Onwuakagba, I.U., Okoye, E.C., Kanu, F.C., Kalu, C.M., Akaeme, D.C., Obaji, O.C., Akosile, C.O. (2023). Population-based stroke risk profile from a West-African Community. *Eneurologicalsci*, 33,100483.
- Osuegbu, O. I., Adeniji, F. O., Ow honda, G. C., Kanee, R. B., Alblihed, M., Batiha, G. E.-

- S., Aigbogun, E. O. (2021). Non-modifiable risk factors (age and sex), stroke types, and outcomes in Rivers State, Nigeria: a retrospective hospital-based study. Preprints, 2021070446. <https://doi.org/10.20944/preprints202107.0446.v1>
- Periyasamy, M., Bharathi, D.P., Vinatha, Devi, M.K. (2023). Typical presentation of malaria with stroke. *J Res Med Dent Sci*, 11 (1), 299-300.
- Rink, C. and Khanna, S. (2011). Significance of brain tissue oxygenation and the arachidonic acid cascade in stroke. *Antioxidants & Redox Signaling*, 14(10), 1889–1903.
- Saengsuwan, J. and Suangpho, P. (2019). Self-perceived and actual risk of further stroke in patients with recurrent stroke or recurrent transient ischemic attack in Thailand. *J Stroke Cerebrovasc Dis*, 28, 632–9. doi: 10.1016/j.jstrokecerebrovasdis.2018.11.001
- Sung, J., Song, Y.M., Choi, Y.H., Ebrahim, S., Davey Smith, G. (2007). Hepatitis B virus seropositivity and the risk of stroke and myocardial infarction. *Stroke*, 38, 1436–1441.
- Talabi, O. (2003). A 3-year review of neurologic admissions in university college hospital Ibadan, Nigeria. *West Afr. J. Med*, 22: 150–151.
- Tamer, Y., Mohammad, H.J., Safi, U.K., Reed, M., Syed, Z.H., Michael, J.B., Ron, B., Salim, S.V., Michelle, C.J., Farhaan, V., Miguel, C., and Khurram, N. (2020). Stroke in young adults: current trends, opportunities for prevention and pathways forward. *Am J Prev Cardiol*, PMID: 34327465. doi: 10.1016/j.ajpc.2020.100085
- Teh, W.L., Abdin, E., Vaingankar, J.A., Seow, E., Sagayadevan, V., Shafie, S., Shahwan, S., Zhang, Y., Chong, S.A., Ng, L.L., Subramaniam, M. (2018). Prevalence of stroke, risk factors, disability and care needs in older adults in Singapore. *BMJ Open*, 8(3), Article e020285, 10.1136/bmjopen-2017-020285
- Thakur, K.T., Lyons, J.L., Smith, B.R., Shinohara, R.T., Mateen, F.J. (2016). Stroke in HIV-infected African Americans: a retrospective cohort study. *J Neurovirol*, 1, 22, 50-55.
- Verhoeven, J.I., Allach, Y., Vaartjes, I.C.H., Klijn, C.J.M., de Leeuw, F.E. (2021). Ambient air pollution and the risk of ischemic and haemorrhagic stroke. *Lancet Planet Health*, 5, e542–e552.
- Warren-Gash, C., Blackburn, R., Whitaker, H., McMenamin, J., Hayward, A. C. (2018). Laboratory-confirmed respiratory infections as triggers for acute myocardial infarction and stroke: a self-controlled case series analysis of national linked datasets from Scotland. *European Respiratory Journal*, 51: 1701794; DOI: 10.1183/13993003.01794-2017
- Wasay, M., Khan, M., Farooq, S., Khowaja, Z.A., Bawa, Z.A., Mansoor Ali, S., Awan, S., Beg, M.A. (2018). Frequency and impact of cerebral infarctions in patients with tuberculous meningitis. *Stroke*, 49, 2288–2293. [Google Scholar]
- Watila, M.M., Nyandaiti, Y.W., Balarabe, S., Bakki, B., Alkali, N.H., Ibrahim, A., Elijah, T., and Chiroma, I. (2019). Aspiration pneumonia in patients with stroke, Northeast Nigeria. *National Journal of Neurology*, 1(3), 48-55. DOI:10.28942/nnj.v1i3.182
- Wellenius, G.A., Burger, M.R., Coull, B.A., Schwartz, J., Suh, H.H., Koutrakis, P., Schlaug, G., Gold, D.R., Mittleman, M.A. (2012). Ambient air pollution and the risk of acute ischemic stroke. *Arch. Intern. Med.*, 172, 229–234.
- WHO Regional Office for Africa (2022). Report on malaria in Nigeria 2022. <https://www.afro.who.int/countries/nigeria/publication/report-malaria-nigeria-2022>
- Zewdie, A., Debebe, F., Kebede, S., Azazh, A., Laytin, A., Pashmforoosh, G., et al. (2018). Prospective assessment of patients with stroke in Tikur Anbessa specialised hospital, Addis Ababa, Ethiopia. *Afr J Emerg Med*, 8(1), 21–4.
- Zhang, F. L., Guo, Z. N., Wu, Y. H., Liu, H. Y., Luo, Y., Sun, M. S. et al. (2017). Prevalence of stroke and associated risk factors: a population based cross sectional study from Northeast China. *BMJ Open*, 7(9):e015758.